

containing a 1,3-dioxolane group: EP-B-0 288 813; EP-B-0 361 272;
EP-B-0 462 156; EP-B-0 351 746




It has been found in accordance with the invention that active-matrix displays in which the ferroelectric smectic phase is stable over a broad temperature range are obtainable by using the compounds of the formula (I). Furthermore, the acute angle is very stable over a broad temperature range, i.e. it is only subject to very small changes. The same applies to the layer leaning angle.

10 In formula (I), X is preferably -OC(=O)-, -OCH₂- or -OC(=O)CH₂CH₂-, particularly preferably -OC(=O).


B¹ is preferably cyclohexane-1,4-diyl, cyclohex-1-ene-1,4-diyl, phenylene-1,4-diyl, unsubstituted, monosubstituted or disubstituted by F, or thiophene-2,5-diyl, particularly preferably cyclohexane-1,4-diyl or thiophene-2,5-diyl.

A¹ is preferably pyrimidine-2,5-diyl (unsubstituted or monosubstituted by F), pyridine-2,5-diyl (unsubstituted or monosubstituted by F), phenylene-1,4-diyl (unsubstituted, monosubstituted or disubstituted by F) or (1,3,4)-thiadiazole-2,5-diyl.


Preferred compounds of the formula (I) correspond to the formulae



12




where R^1 , R^2 are as defined above and



5 is a bivalent radical selected from the group consisting of phenylene-1,4-diyl, unsubstituted, monosubstituted or disubstituted by F, pyrimidine-2,5-diyl, unsubstituted or monosubstituted by F, pyridine-2,5-diyl, unsubstituted or monosubstituted by F, (1,3,4)-thiadiazole-2,5-diyl, indane-2,5-diyl, cyclohexane-1,4-diyl, unsubstituted or monosubstituted by F or CN, cyclohex-1-ene-1,4-diyl, 1,2,3,4-tetra-
10 hydroquinazoline-2,6-diyl




15 is a bivalent radical selected from the group consisting of phenylene-1,4-diyl, unsubstituted, monosubstituted or disubstituted by F, pyrimidine-2,5-diyl, unsubstituted or monosubstituted by F, pyridine-2,5-diyl, unsubstituted or monosubstituted by F, indane-2,5-diyl




20 is a bivalent radical selected from the group consisting of cyclohexane-1,4-diyl, unsubstituted or monosubstituted by F or CN, cyclohex-1-ene-1,4-diyl, (1,3)-dioxane-2,5-diyl, unsubstituted or monosubstituted by CN, thiophene-2,5-diyl, thiophene-2,4-diyl, phenylene-1,4-diyl, unsubstituted, monosubstituted or disubstituted by F,
25 phenylene-1,3-diyl, unsubstituted, monosubstituted or disubstituted by F.

Particularly preferred compounds of the formula (I) correspond to the formulae


13



(Ia1)




(Ib1)



(Ic1)

where:



5


is a bivalent radical selected from the group consisting of phenylene-1,4-diyl, unsubstituted, monosubstituted or disubstituted by F, (1,3,4)-thiadiazol-2,5-diyl, pyrimidine-2,5-diyl, unsubstituted or substituted by F, pyridine-2,5-diyl, unsubstituted or substituted by F ortho to the nitrogen atom, 1,2,3,4-tetrahydroquinazoline-2,6-diyl

10



is a bivalent radical selected from the group consisting of phenylene-1,4-diyl, unsubstituted, monosubstituted or disubstituted by F, pyrimidine-2,5-diyl, unsubstituted or monosubstituted by F, pyridine-2,5-diyl, unsubstituted or monosubstituted by F

15



is a bivalent radical selected from the group consisting of cyclohexane-1,4-diyl, thiophene-2,5-diyl, phenylene-1,4-diyl

20